

AD-A138 514

CURRENT AND TEMPERATURE DATA FROM INSTRUMENT MOORINGS
IN FRAM STRAIT DURING MIZEX 83(U) LAMONT-DOHERTY
GEOLOGICAL OBSERVATORY PALISADES NY K HUNKINS JAN 84
LDGO-84-1 CU-002-97201

1/1

UNCLASSIFIED

F/G 8/3

NL


END

DATE

FILED

3-84

DTIC



MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS 1963-A

AD A138514

CURRENT and TEMPERATURE DATA

FROM INSTRUMENT MOORINGS in FRAM STRAIT

DURING MIZEX 83

Kenneth Hunkins

Technical Report LDGO-84-1

Department of the Navy
Office of Naval Research
Contract No. CU-002-97201

Lamont-Doherty Geological Observatory
of Columbia University
Palisades, New York 10964

Approved for Public Release, Distribution Unlimited

January 1984

ABSTRACT

The purpose of the MIZEX 83 field program in mesoscale oceanography was to monitor mesoscale variability in Fram Strait. An array of instruments was moored in open water near the center of Fram Strait to assist in accomplishing this goal. Current, temperature and pressure measurements from this array are plotted against time. Simple statistics and power spectra for these data are also presented.

| | |
|--------------------|-------------------------------------|
| Accession For | |
| NTIS GRA&I | <input checked="" type="checkbox"/> |
| DTIC TAB | <input type="checkbox"/> |
| Unannounced | <input type="checkbox"/> |
| Justification | |
| By | |
| Distribution/ | |
| Availability Codes | |
| Dist | Avail and/or Special |
| A-1 | |



TABLE OF CONTENTS

| | <u>Page</u> |
|--|-------------|
| Abstract | 1 |
| Contents | 11 |
| Acknowledgements..... | iii |
| Introduction..... | 1 |
| The MIZEX 83 Moorings..... | 3 |
| Oceanographic Setting..... | 5 |
| Mean Currents and Temperatures..... | 7 |
| Time Series..... | 10 |
| Power Spectra..... | 11 |
| References | 12 |
| Figure 1 Location of MIZEX 83 Mooring Array (Bathymetry from Johannessen and Johannessen, 1983)..... | 2 |
| Figure 2 CTD Station 319, 78.5° N, 2.6° E; 25 July 83 (Johannessen and Johannessen, 1983)..... | 6 |
| Figure 3 Mean current vectors at Mooring 4. Depths in meters, 41-day means except for the 158-m record which is a 30-day mean..... | 9 |
| Table 1 Mooring locations and depths..... | 4 |
| Table 2 Mean temperatures, mean current compo- nents and their standard deviation..... | 8 |
| Time Series Plots..... | 13 |
| Power Spectra Plots..... | 53 |

ACKNOWLEDGEMENTS

The MIZEX 83 mooring results represent the combined efforts of many people. The moorings were deployed under the supervision of Ivars Bitte. Jay Ardai, Barry Allen and Peter Bruchhausen assisted with the launching. Recovery was supervised by Jay Ardai with the assistance of Allan Heilscher. The captain and crew of the M/V POLARBJORN provided capable assistance throughout the mooring operations.

Werner Tiemann, Dennis Camp, Barry Allen and Rich Cember assisted with data reduction. Tom Manley wrote a number of the computer programs used in reduction.

The Lamont Arctic Program is funded by the Office of Naval Research under Contract No. CU-002-97201.

Introduction

An array of moored oceanographic recording instruments was placed in Fram Strait at the beginning of the Marginal Ice Zone Experiment in 1983 (MIZEX 83) and recovered at the end of the experiment. The array was situated near the center of the Strait on the sill separating the Arctic Ocean from the Greenland Sea (Figure 1). Three moorings were arranged in a triangle with the fourth mooring at the center of the triangle. Nominal spacing was 10 km, a distance chosen to be close to the Rossby radius of deformation in the region. The array was in open water east of the ice edge.

The objective was to monitor mesoscale oceanographic fluctuations in open water near the ice edge. An oceanographic front is generally associated with the ice edge and two extreme hypotheses might be postulated about mesoscale motion in the area. Such motions could be due to a spectral cascade of energy generated in boundary layers by the mean flow. In this case, spectra in the time domain would be expected to show a monotonic decrease in energy from low to high frequencies, representing a cascade of geostrophic turbulence. An alternate hypothesis would be that instability processes in the mean shear across the front favor the growth of certain discrete frequencies and wavenumbers. In this case the recordings would be expected to show one or more spectral peaks in the sub-inertial frequency range.




Figure 1 - Location of MIZEX 83 Mooring Array. (Bathymetry from Johannessen & Johannessen, 1983).

The MIZEX 83 Moorings

Four subsurface moorings were deployed in water depths of about 2,400 m on June 16 - 17, 1983 from M/V POLARBJORN in an array nominally centered at 78°30' N 002° E (Table 1). The array location was south of the Molloy Deep, where depths exceed 4,000 m, and north of a northwest-southeast trending ridge which rises to a depth of about 1,700 m. Bottom slopes were gentle in the area of the array. All of the moorings and instruments were retrieved 41 days later on July 27, 1983. One of the moorings was then redeployed to record over the next year. It will be retrieved during MIZEX 84.

Aanderaa RCM-5 current meters were used to record current speed and direction, as well as temperature and depth. Manufacturer's specifications for these instruments are:

| | |
|------------------------------|------------------------------|
| Temperature resolution | $\pm 0.01^{\circ} \text{ C}$ |
| Current speed accuracy | $\pm 2 \%$ of actual speed |
| Current direction resolution | $\pm 0.35^{\circ}$ |

All instruments were calibrated before the MIZEX 83 cruise. Although the instruments were equipped with conductivity sensors, range settings were incorrect for this region and no salinity data were obtained. Current meters were suspended at nominal depths of 50 and 150 m below the surface on all four moorings. Additional instruments were suspended at depths of 1,000 below the surface and 100 m above the bottom. Data were sampled every 10 minutes.

The magnetic recording tape failed to transport on four of the twelve instruments deployed. The malfunction was later found to be caused by excessive use of tape head cleaner which did not entirely evaporate before the tape was mounted. This resulted in adhesion of tape to the tape head.

Table 1


MOORING LOCATIONS and DEPTHS

| Mooring Number | Ocean Depth (m) | Latitude N deg min | Longitude E deg min | Aanderaa S.N. | Minimum Depth (m) | Mean Depth (m) | Maximum Depth (m) | Record Length (days) |
|-------------------|-----------------------|-----------------------|------------------------|------------------|-------------------------|----------------------|-------------------------|----------------------------|
| 1 | 2340 | 78 30.42 | 1 57.66 | 5786 | 52 | 55 | 77 | 26.14 |
| | | | | 5461 | 151 | 154 | 179 | 33.63 |
| 2 | 2347 | 78 33.78 | 1 58.80 | 5378 | 61 | 63 | 92 | 41.24 |
| | | | | 6538 | 1015 | 1018 | 1047 | 41.15 |
| 4 | 2439 | 78 36.06 | 1 52.02 | 5785 | 55 | 59 | 106 | 40.78 |
| | | | | 5760 | 158 | 161 | 207 | 40.75 |
| | | | | 6536 | 1014 | 1017 | 1059 | 40.76 |
| | | | | 5889 | 2296 | 2304 | 2309 | 40.74 |

Another instrument was accidentally fouled in the mooring cable during launching so that it was unable to swivel. Temperature data are available for that instrument, but current data are invalid. On three other instruments digital noise, apparently due to poor functioning of the digitization mechanism in these three meters, reduced the length of useable record.

Oceanographic Setting

The moorings were situated in open water 50 km east of the ice edge which trends nearly north and south in this region. An oceanographic profile was made from M/V POLARBJORN by the University of Bergen Oceanographic Group, with a Neil-Brown CTD instrument, just before the moorings were retrieved (Figure 2). Sigma-t was about 27.2 at the surface and increased rapidly with depth in the upper 100 meters so that at 150 m, it was about 27.8. Below 100 m it increased slowly and was still less than 28.0 at 2,000 m. There was a thin layer of warm surface water at about 5° C which extended to a depth of only about 10 m. A thin cold layer at 25 m separated the surface layer from a warmer layer of Atlantic origin with maximum temperatures exceeding 2° C. This layer extended to a depth of 700 m, below which temperatures were negative, closely approaching -1.00° C at 2,000 m. At the surface, salinity was about 34.0 ppt. It increased rapidly with depth in the first 100 m, then increased only slowly at greater depths approaching a value slightly less than 35.0 ppt at 2,000 m.



PROFILE:
 STA: 319 / POS: 78.500°N 2.598°E / TIME: 83-7-25:2.47

Figure 2 - CTD Station 319, 78.5°N, 2.6°E; 25 July 83.
 (Redrawn from Johannessen and Johannessen, 1983).

Mean Currents and Temperatures

Mean values and standard deviations for temperature and current components are presented in Table 2. The shallowest instruments, nominal depth of 50 m, had mean temperatures of 3.62, 2.51 and 2.56° C. The 3.62° C value is from a shorter record and thus cannot be compared directly with the other two means of 2.51 and 2.56° C which are satisfyingly close to each other in value. Their large standard deviations, ± 0.36 and $\pm 0.63^\circ\text{C}$ reflect the fact that these instruments are located in a thermocline, so that any vertical water motions produce large recorded temperature variations.

The two instruments at the 150-m level recorded mean temperatures of 1.67 and 1.62° C, again relatively close in value as are also the two means of -0.77 and -0.74° C at the 1,000-m level. There was only one record from 100 m above the bottom and the mean at that level was -1.00° C, quite close to the temperature at 2,000 m observed for the CTD profile. The standard deviation of $\pm 0.04^\circ\text{C}$ near the bottom indicates the constancy of temperature at that level and also gives some measure of the repeatability of the temperature measurements.

Mean currents at four levels in the water column from near the surface to near the bottom all have a southward component (Figure 3). Mean speeds are relatively high, ranging from 5.2 to 7.6 cm/s. Although there are shears between the different levels, it is perhaps the agreement rather than differences between these means that should be emphasized. The general similarity in speed and direction between the different levels suggests that the mean flow has a large barotropic component. The time series do show some long-period trends extending over several weeks, and means over longer periods

Table 2

MEAN VALUES and STANDARD DEVIATIONS
for TEMPERATURE and CURRENT

| Mooring Number | Instrument Depth (m) | Record Length (days) | \bar{T} (°C) | \bar{u} (cm/s) | \bar{v} (cm/s) |
|-------------------|----------------------------|----------------------------|-------------------|---------------------|---------------------|
| 1 | 52 | 26 | 3.62 ±0.56 | -1.7 ±8.2 | 0.4 ±9.4 |
| | 151 | 29 | 1.67 ±0.30 | 1.6 ±5.3 | -5.0 ±7.3 |
| 2 | 61 | 41 | 2.51 ±0.36 | | |
| | 1015 | 41 | -0.77 ±0.08 | 1.1 ±2.5 | -4.4 ±4.2 |
| 4 | 55 | 41 | 2.56 ±0.63 | -2.0 ±6.6 | -7.5 ±8.2 |
| | 158 | 30 | 1.62 ±0.40 | -2.0 ±5.8 | -6.5 ±7.3 |
| | 1014 | 41 | -0.74 ±0.08 | 0.5 ±2.4 | -5.2 ±3.8 |
| | 2296 | 41 | -1.00 ±0.04 | 1.2 ±3.0 | -7.5 ±3.9 |




Figure 3 - Mean current vectors at Mooring 4. Depths in meters, 41-day means except for the 158-m record which is a 30-day mean.

might have shown different results. The results do indicate that there must have been a large northward flow in another part of the Strait during this period to compensate for the large observed southward transport.

Time Series

The data are presented as plots of temperature, eastward velocity (U) and northward velocity (V) for each instrument. Instrument depth is also plotted. The instruments were all calibrated just before the cruise and the results have been reduced using the corrections indicated by the calibrations. A correction for a magnetic declination of 12.3° W was made to current direction.

Visual examination of the records shows fluctuations of semidiurnal tidal period, as well as longer period events lasting several days which are probably meteorological in origin. A prominent cold feature appears on the shallow temperature record at mooring 4 during July 9 - 10, 1983. Temperature at the 55 m level dropped by nearly 4° C in a few hours and continued cold for two days before returning to its normal level of about $2\frac{1}{2}^\circ$ C in as short a time as it took to drop. The feature is evident but attenuated at 158 m depth on the same mooring. There were no significant currents accompanying this feature. It appears to be a shallow cold water mass which was advected past the array. There were sharp drops in temperature on the shallow instruments at moorings 1 and 2 during the same time period as the more prominent one appeared on mooring 4, but the events on moorings 1 and 2 were of short duration, lasting only a few hours.

Power Spectra

Spectra of kinetic energy for the orthogonal current components generally show an energy decline from lowest to highest frequencies with prominent tidal peaks at semidiurnal and diurnal frequencies. Although some peaks are present at sub-tidal frequencies which are significant at the 95% level, they do not appear consistently on all records. The data thus seem to favor the geostrophic turbulence hypothesis referred to in the introduction. No prominent and consistent peaks of eddy energy are evident. Thus the spectra concur with visual examination of the time series which showed only a single shallow cold eddy in the 40-day period.

Spectral peaks at the semidiurnal peaks, M_2 and S_2 , appear in all spectra with energies generally higher in the north component. This tendency for alignment of tidal motion with the axis of Fram Strait is even more pronounced in the diurnal than in the semidiurnal component. In most of the spectra, the diurnal peak for the north component is only slightly less than the semidiurnal peak. However, the diurnal peak for the east component is small and not even evident in many of the spectra.

REFERENCES

Johannessen, J.A. and O.M. Johannessen, A CTD report from Marginal Ice Zone Experiment (MIZEX) in the Fram Strait region in June - August 1983, Vol. II. Geophysical Institute, Division A; University of Bergen, Bergen, Norway, 5 October 1983.


MOORING 1

RCM-5 No. 5786


52 m Depth

Temperature, Current Components and Depth


Mooring ZI-5786 : Deployed from 16-JUN-1983 to 27-JUL-1983 : Mean depth is 54 meters




Mooring Z1-5786 : Deployed from 16-JUN-1983 to 27-JUL-1983 ; Mean depth is 54 meters



Mooring Z1-5786 : Deployed from 16-JUN-1983 to 27-JUL-1983 : Mean depth is 54 meters



Mooring Z1-5786 : Deployed from 16-JUN-1983 to 27-JUL-1983 : Mean depth is 54 meters




MOORING 1

RCM-5 No. 5461


151 m Depth

Temperature, Current Components and Depth


Mooring Z1-5461 : Deployed from 16-JUN-1983 to 27-JUL-1983 : Mean depth is 151 meters




Mooring Z1-5461 : Deployed from 16-JUN-1983 to 27-JUL-1983 : Mean depth is 151 meters



Mooring Z1-5461 : Deployed from 16-JUN-1983 to 27-JUL-1983 : Mean depth is 151 meters



Mooring Z1-5461 : Deployed from 16-JUN-1983 to 27-JUL-1983 : Mean depth is 151 meters




MOORING 2

RCM-5 No. 5378


61 m Depth

Temperature and Depth

Mooring Z2-5378 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 62 meters



Mooring 22-5378 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 62 meters




MOORING 2

RCM-5 No. 6538


1,015 m Depth

Temperature, Current Components and Depth


Mooring 22-6538 : Deployed from 17-JUN-1983 to 27-JUL-1983 ; Mean depth is 1018 meters




Mooring 22-6538 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 1018 meters



Mooring Z2-6538 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 1018 meters



Mooring Z2-6538 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 1018 meters




MOORING 4

RCM-5 No. 5785


55 m Depth

Temperature, Current Components and Depth


Mooring 24-5785 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 58 meters




Mooring Z4-5785 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 58 meters



Mooring Z4-5785 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 58 meters



Mooring Z4-5785 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 58 meters



MOORING 4


RCM-5 No. 5760

158 m Depth


Temperature, Current Components and Depth

(Depth Record in Three Segments)


Mooring 24-5760 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 162 meters




Mooring Z4-5760 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 162 meters




Mooring 24-S760 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 162 meters




Mooring Z4-5760 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 161 meters



Mooring Z4-5760 : Deployed from 17-JUN-1983 to 27-JUL-1983 ; Mean depth is 165 meters



Mooring Z4-5760 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 158 meters




MOORING 4

RCM-5 No. 6536


1,014 m Depth

Temperature, Current Components and Depth


Mooring Z4-6536 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 1017 meters




Mooring 24-6536 : Deployed from 17-JUN-1983 to 27-JUL-1983 ; Mean depth is 1017 meters



Mooring 24-6536 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 1017 meters



Mooring 24-6536 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 1017 meters




MOORING 4

RCM-5 No. 5889


2,296 m Depth

Temperature, Current Components and Depth


Mooring 24-5889 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 2304 meters




Mooring Z4-5889 : Deployed from 17 JUN-1983 to 27-JUL-1983 : Mean depth is 2304 meters



Mooring 24-5889 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 2304 meters




Mooring 24-5889 : Deployed from 17-JUN-1983 to 27-JUL-1983 : Mean depth is 2304 meters





POWER SPECTRA


of


CURRENT COMPONENTS






































MANDATORY

DISTRIBUTION LIST

1 Director of Defense Research and Engineering
Office of the Secretary of Defense
Washington, DC 20301
Attn: Office, Assistant Director (Research)

Office of Naval Research
Arlington, VA 22217

1 Attn: Code 102-C
1 Attn: Code 200
1 Attn: Code 428 AR
3 Attn: Code 420

6 Director
Naval Research Laboratory
Washington, DC 20375
Attn: Library, Code 2620

1 U.S. Naval Research Laboratory
Code 2627
Washington, DC 20375

1 Office of Naval Research - New York
715 Broadway
New York, NY 10003

12 Defense Documentation Center
Cameron Station
Alexandria, VA 22314

1 Commander
Naval Oceanographic Office
NSTL Station
Bay St. Louis, MS 39522
Attn: Code 02

MIZEX Distribution List

| | | |
|-------------------|----------------------|-------------------|
| E. Augstein | W. J. Campbell | K. Davidson |
| I. Dyer | J. C. Gascard | P. Gudmandsen |
| W. D. Hibler, III | T. Jakobsson | O. M. Johannessen |
| G. A. Maykut | J. Morison | R. H. Shuchman |
| E. C. Monahan | C. Matzler | M. McPhee |
| S. D. Smith | W. O. Smith | P. Wadhams |
| A. Baggeroer | H. Dahlgaard | H. H. Essen |
| B. A. Farrelly | T. Grenfell | J. P. Hollinger |
| K. L. Hunkins | H. Ito | E. Josberger |
| N. Lannelongue | J. Lenz | R. Markson |
| J. Meincke | E. Mollo-Christensen | R. G. Onstott |
| W. Rosenthal | C. Schgounn | G. Sebestyen |
| E. A. Svendsen | G. Symonds | J. K. Vandiver |
| J. C. VanLeer | J. A. Johannessen | D. A. Horn |
| R. Armstrong | A. Heiberg | G. L. Johnson |

1/20/84

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

| REPORT DOCUMENTATION PAGE | | READ INSTRUCTIONS BEFORE COMPLETING FORM |
|---|-------------------------------------|--|
| 1. REPORT NUMBER LDGO-84-1 | 2. GOVT ACCESSION NO. AD-A138514 | 3. RECIPIENT'S CATALOG NUMBER |
| 4. TITLE (and Subtitle) Current and Temperature Data from Instrument Moorings in Fram Strait during MIZEX 83 | | 5. TYPE OF REPORT & PERIOD COVERED |
| | | 6. PERFORMING ORG. REPORT NUMBER |
| 7. AUTHOR(s) Kenneth Hunkins | | 8. CONTRACT OR GRANT NUMBER(s) CU-002-97201 Scope 04 |
| 9. PERFORMING ORGANIZATION NAME AND ADDRESS Lamont-Doherty Geological Observatory of Columbia University Palisades, NY 10964 | | 10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS |
| 11. CONTROLLING OFFICE NAME AND ADDRESS Office of Naval Research Arctic Programs, Code 428 AR Arlington, VA 22217 | | 12. REPORT DATE January 1984 |
| | | 13. NUMBER OF PAGES 69 |
| 14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office) | | 15. SECURITY CLASS. (of this report) Unclassified |
| | | 15a. DECLASSIFICATION/DOWNGRADING SCHEDULE |
| 16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited. | | |
| 17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) | | |
| 18. SUPPLEMENTARY NOTES | | |
| 19. KEY WORDS (Continue on reverse side if necessary and identify by block number) MIZEX 83, moorings, mesoscale variability, Fram Strait | | |
| 20. ABSTRACT (Continue on reverse side if necessary and identify by block number) The purpose of the MIZEX 83 field program in mesoscale oceanography was to monitor mesoscale variability in Fram Strait. An array of instruments was moored in open water near the center of Fram Strait to assist in accomplishing this goal. Current, temperature and pressure measurements from this array are plotted against time. Simple statistics and power spectra for these data are also presented. | | |

DD FORM 1 JAN 73 1473

EDITION OF 1 NOV 68 IS OBSOLETE
S/N 0102-LF-014-6601

Unclassified

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DA
FILM